



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The finite discontinuities occur in both cases for the parameters $\alpha, \beta = \pm 1$.

Also solved by M. E. Graber, G. W. Greenwood, F. D. Posey, S. A. Corey, G. B. M. Zerr, and J. O. Mahoney.

MECHANICS.

170. Proposed by M. E. GRABER, A. M., Heidelberg University, Tiffin, Ohio.

Prove that the moment of inertia of an ogival head rotating about its geometrical axis is $\frac{\pi w}{g} \int_0^{R\sqrt{4n-1}} y^4 dx$, where w is the weight in pounds of a cubic foot of material, R the radius of the base of the ogive, and n the diameter of projectile.

Solution by LYTLE BROWN, Lehigh University, and the PROPOSER.

An ogival head is one-half the solid generated by the revolution of a segment of a circle about its chord. The equation of the generating curve is $y = \sqrt{(4n^2 R^2 - x^2)} - (2n-1)R$, the origin being the center of the base of the ogive. If $y=0$, the resulting value of x is the length of the head, that is, the length of the head is $R\sqrt{4n-1}$. The moment of inertia of an elementary cylinder about the axis of x is $4\pi y^2 \frac{\omega}{g} \cdot y^2 dx$, or $dI = \frac{\pi \omega}{2g} y^4 dx$.

$$\therefore I = \frac{\pi \omega}{g} \int_0^{R\sqrt{4n-1}} y^4 dx.$$

AVERAGE AND PROBABILITY.

155. Proposed by E. B. WILSON, Ph. D., Yale University.

The game of craps is played with two dice. If the player throws 7 or 11 on the first throw he wins. If he throws 12, 2, or 3 he loses. If the player throws any other number, that is to say, 4, 5, 6, 8, 9, 10, he is obliged to continue throwing until he throws that number again or until he throws 7. If he succeeds in throwing his first throw before he does 7, he wins—otherwise he loses. Required the odds against him. (Note that he can continue throwing indefinitely without getting either his original throw or the 7).

REMARK. A correct solution of this problem is given by Dr. Zerr, in Vol. X, No. 3, p. 81. Mr. J. E. Sanders sent in a different solution, his answer being $\frac{1}{11}$, in favor of the player.

Problem 131 should be numbered 158. Three solutions of this problem have been received, none of which agree.

Problem 157 should be numbered 159.

MISCELLANEOUS.

144. Proposed by IRA M. DeLONG, Professor of Mathematics in the University of Colorado, Boulder, Col.

Determine the number of distinct spherical polygons of n sides formed by arcs of n given great circles on a sphere, each arc to be greater than 0 degrees, and less than 360 degrees, and no two sides of any polygon to lie on the same circle.